

[psng]mm  
 [1] 1 [1] 1 [1] 1 textbf{cmbxti10} textbf{ss cmbssbx10} mathbf{cmbxti10} mathbf{ss cmbssbx10} upmath  
 eurm10 AMSa msam10 0upmath19 0upmath16 0upmath40 3AMSa36 3AMSa3E  
 normalcmrmit boldcmrbxit [1] 1 [1] 1 [1] 1 normalcmrbxit boldcmrbxit normalcmssbxn bold-  
 cmssbxn @mathgroup@group @mathgroup@normal@groupeurmn @mathgroup@bold@groupeurn @math-  
 group@group @mathgroup@normal@groupmsamn @mathgroup@bold@groupmsamn @group19 @group16  
 @group40 @group36 @group3E  
 [1] 1 [1] 1 [1] 1 OT1cmrbxit boldOT1cmrbxit OT1cmssbxn boldOT1cmssbxn UPMUeurmn UPM-  
 boldUeurbn AMSaUmsamn  $\Gamma 0UPM^{19}\Gamma 0UPM^{16}\Gamma 0UPM^{40}\Lambda 3AMSa^{36}\Lambda 3AMSa^{3E}$   
 [Triaxial stellar systems following the  $r^{1/n}$  luminosity law]Triaxial stellar systems following the  $r^{1/n}$   
 luminosity law: an analytical mass–density expression, gravitational torques and the bulge/disc interplay  
 [I. Trujillo et al.] I. Trujillo<sup>1</sup>itc@lliac.es, A. Asensio Ramos<sup>1</sup>, J. A. Rubiño-Martín<sup>1</sup>, Alister W.  
 Graham<sup>1,2</sup>,

Accepted 0000 December 00. Received 0000 December 00; in original form 0000 October 00

firstpage–lastpage 2001

document

firstpage

abstract

We have investigated the structural and dynamical properties of triaxial stellar systems whose surface brightness profiles follow the  $r^{1/n}$  luminosity law – extending the analysis of Ciotti (1991) who explored the properties of spherical  $r^{1/n}$  systems. A new analytical expression that accurately reproduces the spatial (i.e. deprojected) luminosity density profiles (error less than 0.1%) is presented for detailed modelling of the Sérsic family of luminosity profiles. We evaluate both the symmetric and the non-axisymmetric components of the gravitational potential and force and compute the torques as a function of position. *For a given triaxiality, stellar systems with smaller values of  $n$  have a greater non-axisymmetric gravitational field component.* We also explore the strength of the non-axisymmetric forces produced by bulges with differing  $n$  and triaxiality on systems having a range of bulge–to–disc ratios. The increasing disc–to–bulge ratio with increasing galaxy type (decreasing  $n$ ) is found to heavily reduce the amplitude of the non-axisymmetric terms, and therefore reduce the possibility that triaxial bulges in late-type systems may be the mechanism or perturbation for non-symmetric structures in the disc.

Using seeing-convolved  $r^{1/n}$ –bulge plus exponential–disc fits to the K-band data from a sample of 80 nearby disc galaxies, we probe the relations between galaxy type, Sérsic index  $n$  and the bulge–to–disc luminosity ratio. These relations are shown to be primarily a consequence of the relation between  $n$  and the total bulge luminosity. In the K-band, the trend of decreasing bulge–to–disc luminosity ratio along the spiral Hubble sequence is predominantly, although not entirely, a consequence of the change in the total bulge luminosity; the trend between the total disc luminosity and Hubble type is much weaker.













